

Overview of functionality in the ONETEP code

Nicholas D.M. Hine

Assistant Professor, University of Warwick

ONETEP Masterclass 2015, Cambridge, 15th August 2015



ONETEP

Overview


- LR-TDDFT
- Implicit Solvent
- van der Waals
- Metallic systems
- External Pressure
- PAW
- Hybrid Functionals
- Molecular Dynamics
- Transport
- Population Analysis
- Strong Correlations
- Constrained DFT
- Phonons
- Future Plans

Linear-Response TDDFT

Capability: Calculate localised excitations and optical spectra

People: Tim Zuehlsdorff, Jian-Hao Li, Peter Haynes, Nicholas Hine

- Linear-scaling time-dependent density-functional theory in the linear response formalism, T. J. Zuehlsdorff, N. D. M. Hine, J. S. Spencer, N. M. Harrison, D. J. Riley, and P. D. Haynes, *J. Chem. Phys.* 139, 064104 (2013).
- Linear-scaling time-dependent density-functional theory (TDDFT) beyond the Tamm-Dancoff approximation: obtaining efficiency and accuracy with in situ optimised local orbitals, T. J. Zuehlsdorff, N. D. M. Hine, M.C. Payne, and P. D. Haynes (2015).



Quantified Natural Transition Orbitals will be available soon:


- Identifying and Tracing potential energy surfaces of electronic excitations via their transition origins: application to Oxirane, J.-H. Li, T.J. Zuehlsdorff, M. C. Payne, N. D. M. Hine, *Phys. Chem. Chem. Phys.* 17, 12065 (2015)

Implicit Solvent

Capability: Simulate molecules, nanostructures etc in a dielectric medium

People: Jacek Dziedzic, Arash Mostofi, Chris Skylaris

- Minimal parameter implicit solvent model for ab initio electronic structure calculations, J. Dziedzic, H. H. Helal, C.-K. Skylaris, A. A. Mostofi, and M. C. Payne. *Europhysics Letters* 95, 43001 (2011).
- Large-scale DFT calculations in implicit solvent - a case study on the T4 lysozyme L99A/M102Q protein, J. Dziedzic, S. J. Fox, T. Fox, C. S. Tautermann, and C.-K. Skylaris, *Int. J. Quantum Chem* 113, 771, (2013)




Metallic Systems

Capability: Treatment of large metallic systems eg metallic nanoparticles (Asymptotic Cubic-Scaling, but near-linear for medium/large systems)

People: Alvaro Ruiz-Serrano, Chris Skylaris

- A variational method for density functional theory calculations on metallic systems with thousands of atoms, A. Ruiz-Serrano and C.-K. Skylaris, J. Chem. Phys. 139, 054107 (2013).




van der Waals

Capability: Fully non-local van der Waals Density Functionals, using functionals eg Dion et al, and fast method of Roman-Perez & Soler.

People: Lampros Andrinopoulos, Gabriel Constantinescu, Fabiano Corsetti, Nicholas Hine, Arash Mostofi

- Article in Preparation: L. Andrinopoulos, F. Corsetti, N. Hine, A. Mostofi
- Used in: Energy landscape and band-structure tuning in realistic MoS₂/MoSe₂ heterostructures G. C. Constantinescu, N. D. M. Hine Phys. Rev. B 91, 195416 (2015)

$$E_c^{\text{nl}} = \frac{1}{2} \int \int d\mathbf{r} d\mathbf{r}' \rho(\mathbf{r}) \phi(\mathbf{r}, \mathbf{r}') \rho(\mathbf{r}') \Rightarrow E_c^{\text{nl}} = \frac{1}{2} \int \int d\mathbf{r} d\mathbf{r}' \rho(\mathbf{r}) \phi(q, q', r) \rho(\mathbf{r}')$$




Update: Now have VDW-DF1, VDW-DF2, OPTPBE, OPTB88, C09, etc

External Pressure

Capability: Simulated systems under external pressure, via enthalpy method

People: Niccolo Corsini, Nicholas Hine, Peter Haynes

- Simulations of Nanocrystals Under Pressure: Combining Electronic Enthalpy and Linear-Scaling Density-Functional Theory, N. R. C. Corsini, A. Greco, N. D. M. Hine, C. Molteni, P. D. Haynes, J. Chem. Phys., 139, 084117 (2013).
- *Pressure-induced amorphisation and a new high density amorphous metallic phase in matrix-free Ge nanoparticles*, N. R. Corsini, [...], N.D.M. Hine, P. D. Haynes, C. Molteni, A. Sapelkin (under review) 2015.




Projector Augmented Wave

Capability: Use PAW treatment of core electrons rather than norm-conserving pseudos. Effectively an all-electron treatment, enabling accurate calculations on high-Z elements & transition metals.

People: Nicholas Hine

- Article in Preparation: N. D. M. Hine, *Linear-Scaling Plane-Wave Density Functional Theory using the Projector Augmented Wave Method*
- Used in: Energy landscape and band-structure tuning in realistic MoS₂/MoSe₂ heterostructures G. C. Constantinescu, N. D. M. Hine Phys. Rev. B 91, 195416 (2015)




Hybrid Functionals

Capability: Functionals including exact exchange (eg B3LYP)

People: Jacek Dzedzic, Chris Skylaris

- Linear-Scaling calculation of Hartree-Fock exchange energy with Non-Orthogonal Generalised Wannier functions, J. Dzedzic, Q. Hill, C. K. Skylaris, J. Chem. Phys., 139, 214103 (2013)





Molecular Dynamics

Capability: Perform Dynamics on large systems, with density matrix extrapolation

People: Valerio Vitale, Simon Dubois, Chris Skylaris

- Anharmonic infrared spectroscopy through the Fourier transform of time correlation function formalism in ONETEP, V. Vitale, J. Dziedzic, S. M. M. Dubois, H. Fangohr, C.-K. Skylaris, *J. Chem. Theor. Comput.* 11, 3321 (2015).
- Acid-base dissociation mechanisms and energetics at the silica–water interface: An activationless process, B. M. Lowe, , C.-K. Skylaris, N. G. Green, *Journal of Colloid and Interface Science* 451, 231 (2015).



$$1: \mathbf{v}' = \mathbf{v}_n + \frac{\Delta t}{2m} * \mathbf{F}_n \quad (2)$$

$$2: \mathbf{r}_{n+1} = \mathbf{r}_n + \Delta t * \mathbf{v}' \quad (3)$$

$$3: \text{Compute ionic forces } \mathbf{F}_{n+1} \quad (4)$$



$$4: \mathbf{v}_{n+1} = \mathbf{v}' + \frac{\Delta t}{2m} * \mathbf{F}_{n+1} \quad (5)$$

Transport

Capability: Perform Dynamics on large systems, with density matrix extrapolation

People: Simon Dubois, Rob Bell, Arash Mostofi

- Electronic transport calculations in the onetep code: Implementation and applications, R. A. Bell, S.M.M. Dubois, M. C. Payne, A. A. Mostofi, *Comput. Phys. Commun.* 193, 78 (2015).
- *Improving the conductance of carbon nanotube networks through resonant momentum exchange*, R. A. Bell, M. C. Payne, A. A. Mostofi, *Phys. Rev. B* 89, 245426 (2015).




Population Analysis

Capability: Natural Population Analysis, Hirschfeld, DDEC, etc, for forcefield fitting and electrostatics

People: Louis Lee, Danny Cole, Chris Skylaris

- Polarized Protein-Specific Charges from Atoms-in-Molecule Electron Density Partitioning, L. P. Lee , D. J. Cole , C.-K. Skylaris, W. L. Jorgensen , and M. C. Payne, J. Chem. Theory Comput. 9, 2981 (2013)
- Natural bond orbital analysis in the ONETEP code: Applications to large protein systems, L. P. Lee, D. J. Cole, M. C. Payne, and C.-K. Skylaris, J. Comp.Chem. 34, 429 (2013).




Strong Correlations

Capability: DFT+U (penalise non-integer occupations); Dynamical Mean Field Theory

People: David O'Regan, Cedric Weber, Nicholas Hine, Arash Mostofi

- Renormalization of myoglobin–ligand binding energetics by quantum many-body effects, C. Weber, D. J. Cole, D. D. O'Regan, M. C. Payne, PNAS 111, 5790 (2014).
- Importance of many body effects in the kernel of hemoglobin for ligand binding, C. Weber, D. D. O'Regan, N. D. M. Hine, P. B. Littlewood, G. Kotliar and M. C. Payne, Physical Review Letters, 110, 106402 (2013);
- Vanadium dioxide : A Peierls-Mott insulator stable against disorder, C. Weber, D. D. O'Regan, N. D. M. Hine, M. C. Payne, G. Kotliar and P. B. Littlewood, Physical Review Letters 108, 256402 (2012);
- Ligand Discrimination in Myoglobin from Linear-Scaling DFT+U, D. J. Cole, D. D. O'Regan, and M. C. Payne, J. Phys. Chem. Lett. 3, 1448 (2012);
- Linear-scaling DFT+U with full local orbital optimization, D. D. O'Regan, N. D. M. Hine, M. C. Payne and A. A. Mostofi, Phys. Rev. B 85, 085107 (2012).




Constrained DFT

Capability: Constrain populations & spins of certain regions of a system (eg for studying charge transfer reactions)

People: Gilberto Teobaldi David O'Regan, Nicholas Hine, Arash Mostofi

- Article(s) in Preparation



Phonons

Capability: Calculate vibrational frequencies, free energies, phonon modes etc, via a) finite-difference approach, b) linear-response methods

People: Fabiano Corsetti, Gabriel Constantinescu

- Article(s) in Preparation

$$E^{\text{tot}} = E^{\text{eq}} + \frac{1}{2} \sum_{a,\alpha,\kappa,a',\alpha',\kappa'} u_{a,\alpha,\kappa} \phi_{\kappa,\kappa'}^{\alpha,\alpha'}(a,a') u_{a',\alpha',\kappa'},$$

$$\phi_{\kappa,\kappa'}^{\alpha,\alpha'}(a,a') = \frac{\partial^2 E}{\partial u_{a,\alpha,\kappa} \partial u_{a',\alpha',\kappa'}}.$$

$$D_{\kappa,\kappa'}^{\alpha,\alpha'}(\mathbf{q}) = \frac{1}{\sqrt{M_\alpha M_{\alpha'}}} \sum_a \phi_{\kappa,\kappa'}^{\alpha,\alpha'}(a,0) e^{-i\mathbf{q}\cdot\mathbf{R}_a},$$

Future Plans

- TD-hybrid Functional
- Linear Response Phonons
- Spin Orbit
- Many-body Methods

Conclusions

- Many exciting new developments!
- **Long-term vision:** a simulation toolset enabling computational discovery with true predictive power for nanomaterials.

THE UNIVERSITY OF
WARWICK



Centre for Scientific Computing

EPSRC
Engineering and Physical Sciences
Research Council



The Leonardo Trust



THE WINTON PROGRAMME FOR THE
Physics of Sustainability

Imperial College
London

UNIVERSITY OF
Southampton



UNIVERSITY OF
CAMBRIDGE